

## **DIVERSITY OF AVIFAUNA OF KUKKARAHALLI LAKE AND MANASA GANGOTRI UNIVERSITY CAMPUS: MYSORE CITY, KARNATAKA, INDIA**

**Suchitra G\***

Department of Zoology, Maharani's Science College for Women, Mysore- 570 005

---

### **ABSTRACT**

Avifauna encompasses a wide range of bird species, each with its unique characteristics, behaviors, and adaptions. The field study was conducted to examine the diversity of birds, status, occurrence and abundance of avifauna in Mysore city, Karnataka during March 2015 to July 2016. Sampling was done byline transect method and local visual count method, collected data from 2 study area were subjected to estimate diversity of species. A total 73 species under orders were recorded. Order Passeriformes was found to be most dominant with 35 species followed by Apodiformes, Galliformes, Podicipediformes, Psittaciformes, Strigiformes, with 1 species respectively.

### **Keywords:**

Avifaunal Diversity, Line transect method, Residential status,Mysore city.

---

### **INTRODUCTION**

Birds are bipedal, warm-blooded feathered creatures (Ayyar, 1964). Known for their ecological, economical, ethical, medicinal and scientific values (Ali S and Ripley 1996). They have a worldwide distribution, living in and around oceans, rivers, forest and mountains. They are the most noticeable group in the animal kingdom (Lameed, 2011). They inhabit all the ecosystem across the globe. There has been a wide range in size of birds that is from Bee humming 5cm to Ostrich 2.5m. Birds are some of the most prominent species of the Earth's biodiversity and being sensitive to environmental changes (main streaming). Birds are social animals that communicate with visual signs, calls and songs. They display social behaviors such as cooperative breeding and hunting, flocking and mobbing of predators (Lameed, 2011). Their behaviour patterns and reproductive ability have most often been used to examine the long term affects of habitat fragmentation. Hence, they are the good indicators of ecological status of any given ecosystem (Bilgrami, 1995). The ornithological information is utilized to show the impacts of natural changes on biodiversity, currently plentiful endeavors are being made in a few far-off regions (Hussainet al., 2011). They have been used to evaluate the environment throughout the history as 'bio monitors' and; the changes in their population (Bilgrami, 1995). Birds detect changes in the environment which cannot be detected or observed by physical parameters. Birds are also good biological control. They consume insect such as mosquitoes, beetles and stem borers which are pests (Ezealor, 2001). Birds have beneficial interactions with forest plants. The beneficial interactions include pollination and seed dispersal. Flowers of some plant species have been discovered to be visited by birds (Anderson et al., 2006). Birds play a noteworthy function in ecosystem. Birds are an imperative component of biodiversity and their occurrence and distribution are an important phenomenon to understand the overall picture of habitat (Chauhan et al., 2008). Birds have a good system for spreading plant seeds that makes them agents of dispersal. They eat fruits and swallow the seeds of plants. When they dispose of their waste, the seeds are disposed along with it. Birds droppings (faeces) provide good fertility to the soil upon which they are dropped, giving the seeds very good conditions with which to grow (Anderson et al., 2006). Scavenger birds, such as the pied crow (*Corvus albus*), contribute to biomass recycling and to some degree reduce levels of disposal wastes. Frugivorous birds play an important role in seed dispersal of fleshy fruit producing plants (Stevenson and Fanshawe, 2002). Birds are also important in plant pollination as demonstrated by sun-birds, which participate in crossbreeding of flowering plants, especially those with bird-pollination syndrome (Judd et al., 2008). The ecosystem services are important for many communities, and to ensure that birds can fulfil these biological roles at an appropriate level for current and future generations, there is a pressing need to study the dynamics and socioeconomics of bird diversity outside protected areas, especially in urban areas (Gatesire et al., 2014). Portraying and clarifying spatial examples in species variety are essential

strides in moderating worldwide biodiversity (Lee *et al.*, 2004). As the quantity of bird species possessing different altitudinal belts or, life zones' (Ali S ,2002). Rely upon climatic changes joined by comparing changes in vegetation (Sekercioglu and Cagan Hakki 2006).

The present study was under taken to assess the species diversity, abundance, status and distribution of avifauna in different habitats within Mysore city, Karnataka, India.

## MATERIALS AND METHODOLOGY

### Study area

For the research study two places were selected in the Mysore city

I. Site- 1 (Kukkarahalli lake)

Coordinates: 12.3098° N ,76.6326° E

Kukkarahalli lake was built in 1864, to provide drinking water to Mysore city, and came under the custody of the university of Mysore city in 1960. The original catchment area of the lake was~4.5 sq km. And the lake is situated 3km from Mysore city. It is adjacent to the University of Mysore and Kalamandir. The lake is large with an area of 150 acres, receives both south-western and the north-east monsoons with an average rainfall 782mm. This lake drains a catchment area of more than 414 square kilometres (160 sq. mi) and the water body spread over 62 hectares with the maximum depth 5m(16ft).

II. Site-2 (Manasa Gangotri campus)

Coordinates: 12.30210° N, 76.6500° E

The University of Mysore is a public state university in Mysore, Karnataka, India. The university was founded during the region of Krishnaraja Wodeyar IV, the Maharaja of Mysore. It opened on 27 July 1916. The Manasa Gangotri campus is confined within the area of 739 acres of picturesque land containing a sprawling Kukkarahalli lake surrounded by verdurous tress extending to an area of 261 acres, The campus has rich presence of flora which provide wide range of habitats for the birds. Also, some of areas within in the university is quite silent which indirectly promote the avian diversity.

### Methods

The field study observations were conducted twice a month from March 2015 to July 2016 for a period of 5 months to record the avifauna diversity. Birds are sighted during their peak activity from 6:30 am to 8:30 am in the morning and 5:00 pm to 6:30 pm in the evening. For the interpretation of collected data the year was divided into three seasons (i) Spring- March, (ii) Summer- April and May, (iii) Early monsoon-June and July.

The study area was surveyed for recording avifauna diversity by applying line transect method, Local visual method.

The birds were identified using 10×50 DPS I Field 6.5° Olympus binoculars, Photography was done by making use of Cannon HD 30X Optimal Zoom Camera. The recorded birds were identified based on their morphological features such as beak shape, colour, type of foot (e.g., webbed or non-webbed), colour of shank, foot and feather colour with the help of field guide and various key books (Salim Ali, 2002, Grimmet and Inskip, 2007)

The check list of species was prepared following Salim Ali (2002) and (Grimmet and Inskip, 2007)

Order	Family	Common names	Scientific names	Residenti al status	Abundan ce	Feedin G Guild
Passeriformes	Oriolidae	Eurasian golden oriole	<i>Oriolus oriolus</i>	RM	C	OV
	Dicruidae	Black drango	<i>Dicrurus macrocercus</i>	R	VC	IV
	Pycnonotidae	Red-whiskeredbulbul	Pycnonotidae	R	VC	FV
		Red vented bulbul	<i>Pycnonotuscafer</i>	R	C	GV
		White-eared bulbul	<i>Pycnonotidae leucotis</i>	R	VC	GV
	Nectariniidae	Purple rumped sunbird	<i>Nectarinia zeylonica</i>	R	VC	IV
		Loten's sunbird	<i>Nectarinia lotenia</i>	R	VC	IV
		Purple sunbird	<i>Nectarinia astiatica</i>	R	C	NV
		Small sunbird	<i>Nectarinia minima</i>	R	VC	NV
	Muscicapidae	Oriental magpie robin	<i>Copsychus saularis</i>	R	C	IV
		Pied bush chat	<i>Saxicola caprata</i>	R	C	IV
	Cisticolidae	Ashy prinia	<i>Prinia socialis</i>	R	C	IV
		Common tailor bird	<i>Orthotomus sutorius</i>	R	C	IV
	Timaliidae	Red capped babbler	<i>Timala pileata</i>	R	C	IV
		Yellow-breasted babbler	<i>Macronous gularis</i>	R	C	IV

	Motacillidae	Large pied wagtail	Motacilla maderaspatensis	R	VC	IV
	Corvidae	House crow	Corvus splendens	R	VC	OV
		Jungle crow	Corvus macrorhynchos	R	VC	OV
	Paridae	Great tit	Parus major	R	C	IV
		Pied tit	Parus nuchalis	R	C	IV
	Sturnidae	Common myna	Acridotheres tristis	R	C	OV
		Jungle myna	Acridotheres fuscus	R	UC	OV
	Aegithinidae	Common Iora	Aegithinatiphia	R	UC	IV
	Leiothrichidae	Large greybabbler	Turdoides malcolmi	R	UC	GV
		Jungle babbler	Turdoides striatus	R	C	GV
	Passeridae	House sparrow	Passer domesticus	R	O	CV
Pelecaniformes	Ardeidae	Grey heron	Ardea cinerea	RM	O	CV
		Black- crowned night- heron	Nycticorax nycticorax	R	O	CV
		Large egret	Casmerodius albus	RM	UC	CV
		Indian pond-heron	Ardeola grayii	R	VC	CV
		Little egret	Egretta garzetta	R	C	CV
		Cattle egret	Bubulcus ibis	RM	C	CV

		Median egret	Ardea intermedia	RM	O	CV
	Pelecanidae	Spot-billedpelican	<i>Pelecanus philippensis</i>	RM	UC	
	Threskiornithidae	Oriental whiteibis	<i>Threskiornis melanocephalus</i>	R	UC	OV
		Glossy ibis	<i>Plegadis falcinellus</i>	RM	O	PV
		Black ibis	<i>Pseudibis papillosa</i>	R	R	IV
Coraciformes	Meropidae	Blue-cheekedbee-eater	<i>Merops persicus</i>	RM	R	IV
		Blue-tailedbee-eater	<i>Merops philippinus</i>	RM	R	IV
	Alcedinidae	White- breasted kingfisher	<i>Halcyon smyrnensis</i>	R	VC	PV
		Small bluekingfisher	<i>Alcedoatthis</i>	RM	VC	IV
Colbiformes	Columbidae	Spotted dove	<i>Spilopelia chinensis</i>	R	C	GV
		Blue rockpigeon	<i>Columbalivia</i>	R	C	GV
		Little browndove	<i>Streptopelia senegalensis</i>	R	C	OV
Galliformes	Phasianidae	Indian peafowl	<i>Pavo cristatus</i>	R	VC	OV
Cuculiformes	Cuculidae	Asiankoel (Female and Male)	<i>Eudynamys scolopaceus</i>	R	C	FV
		Greatercoucal	<i>Centropus sinensis</i>	R	C	IV
		Lesser coucal	<i>Centropus bengalensis</i>	R	C	IV

Charadriformes	Jacanidae	Bronze winged jacana	Metopidiusindicus	R	C	IV
	Charadridae	Red-wattledlapwing	Vanellus indicus	R	VC	IV
Accipitriformes	Accipitridae	Brahminy kite	Haliasturindus	R	C	CV
		Black kite	Milvus migrans	R	C	CV
		Shikra	Accipiterbadius	R	C	CV
		Long-leggedbuzard	Buteno rufinus	R	C	CV
Gruiformes	Rallidae	Commomcoot	Fulica atra	RM	C	IV
		Purple moorhen	Porphyrio porphrio	R	C	IV
		Commomoorhen	Gallinula chloropus	RM	O	IV
		White- breasted waterhen	Amaurornis phoenicurus	R	O	IV
Piciformes	Megalaimidae	Coppersmithbarbet	Megalaima haemacephala	R	UC	FV
	Picidae	Lesser golden-backed woodpecker	Dinopium benghalense	R	R	IV
		Commongolden-backed woodpecker	Dinopium javanense	R	R	IV
Suliformes	Phalacrocoracidae	Great cormorant	Phalacrocorax carbo	RM	O	PV
		Little cormorant	Microcarboniger	RM	O	PV

		Darter	Anhingidae	RM	O	PV
Psittaciformes	Psittaculidae	Rose-ringedparakeet	<i>Psittacula krameri</i>	R	VC	FV
Apodiformes	Apodidae	Asian palm-swift	<i>Cypsiurus balasiensis</i>	R	C	IV
Strigiformes	Strigidae	Spotted owlet	<i>Athene brama</i>	R	VC	CV
		Asian openbill stork	<i>Anastomus oscitans</i>	R	O	CV
Podicipediformes	Podicipedidae	Little grebe	<i>Tachybaptus ruficollis</i>	R	VC	

R- rare, C- common, VC- very common, O- omnivores, IV- Insectivores, PV piscivores,




Figure 1: Representing occurrence of order with respect to belonging species of birds.

### Result and Discussions

A total of 73 species of birds belonging to families were recorded from the two study areas of Mysore city. On the basis of the order wise species occurrence the order Passeriformes dominated with  $n=27$  followed by the Pelecaniformes, Apodiformes, Galliformes, podicipediformes, Psittaciformes, Strigiformes with  $n=1$  were represented from total collected species (Fig:1). In the present study it is noticed that order Passeriformes was found to be predominated from other 7 orders. Similar results of abundance were also reported by several ornithological studies from India.

Among the 2 study areas. Kukkarahalli lake is observed with more bird species, while Manasagangotri was found to be least. The abundance calculated varied among the species recorded. Based on the percentage of relative abundance it was revealed that 4 species of birds were very common.

The seasonal distribution pattern showed two peaks of species richness, Shannon diversity, equitability, and evenness index, one in early winter and the other in summer. Depending on seasonality, many species breed in late winter, which contributes to more nesting and less roaming in late winter and hence low sighting was revealed. On the contrary, in the spring probably eggs hatch and birds can be seen roaming as they gather food for the new born. New individuals can also contribute to high richness and diversity of spring.

Flowering in the early winter assures food availability and it could also be an important cause for high species richness, while comparatively low vegetation thickness in spring and summer can also contribute to high species richness and diversity mainly due to the fact that as there is more exposure, birds can be easily sighted. Low sighting of the birds in monsoon could be due to the less activity (Anand *et al.* 2007). The daylight is very poor; sometimes the entire area may be occupied by clouds resulting into poor visibility in monsoon. Chilling winds worsen the wet conditions. These result into low species distribution and low sightings in the study area (Anand *et al.* 2007). The species still remain in the study area are habitat specific ones, which can tolerate the hostile conditions and some species become secretive to reside in the dense canopy and thus can not be sighted in the transects. Most of the species encountered in transects during monsoon breed in this season. These conditions also result into meager food availability. It is evident from the study, that frugivorous and insectivorous birds (52%) constitute majority of the bird community in the study area.

Therefore, the food availability for these birds is extremely scanty during the monsoon season. This is another reason for their less sighting and low species richness and diversity. Subsequent seasons show better food availability due to increased sunlight and temperature as well as reduction in rainfall. Insect population rises from October onwards and hence the birds start coming back. The species overlapping pattern starts getting restored. High  $\alpha$ - and  $\beta$ -diversity during spring and summer is attributed to the availability of diverse food.

Majority of the flora of study area shows flowering and fruiting during this period. The study revealed the presence of large number of nectarivorous birds (sun birds), though their number was large but the species are few in number. As per the present study, mixed moist deciduous, and dry deciduous forests in the study area are best habitats for the birds as far as the number and diversity is concerned. As the most serious loss of the biodiversity value occurs in the transformation of original landscapes to croplands due to human interference (Pramod *et al.* 1997), evaluation of bird communities from various study sites from the Western Ghats is essential for planning “biodiversity-friendly” developmental activities.

## REFERENCES

1. Ali S 2002 The Book of Indian Birds (13<sup>th</sup> Revised Edition). Oxford University Press, New Delhi.
2. Ali S. 1949 Bird friends and foes of cultivator. Indian Farming.
3. Ali S. 1971 Sunder Lal Hora memorial lecture. Ornithology in India: its past, present and future. Proc Indian Nat Sci Acad.
4. Anderson SH, Dave K, Alastair W. ROBERTSON3 , Jenny J. LADLEY2 , John G 2006 Birds as pollinators and Dispersers: a casestudy from New Zealand. *Acta Zoologica Sinica*. 52(Supplement): 112–115, 2006
5. Balachandran, S., A.R. Rahmani, N. Ezhilarasi, S. Babu, J.P.P. Chakravarthy & T. Ramesh (2005). Revaluation of bird community structure of Palni Hills with special reference to threatened and endemic species. Final Report. Bombay Natural History Society, Mumbai, 105pp
6. Bilgrami KS (1995). Concept and Conservation of Biodiversity. CBS Publishers and Distributors, Delhi.
7. Robertson H A and Hackwell K R. 1995 Habitat preferences of birds in seral kahikatea. *Dacrycarpus dacrydioides*(podocarpaceae) forest of south wetland, New Zealand. *Bio con* 71:275-280
8. Chauhan, R.R., Shingadia H.U. & Sakthivel V. (2008). Survey of avifauna of Borivali mangroves along the coast of Mumbai. *Nature Environmental and Pollution Technology* 7(2): 229–233.
9. Ezalor AU. 2001. **Important Bird Areas in Africa and Associated Islands: Priority Sites for Conservation ... Pisces Publications**; Cambridge: Birdlife International,
10. Gibbs, J.P. 1993, Importance of small wetlands for the persistence of local populations of wetland- associated animals. *Wetlands* 13, 25–31
11. Grimmett, R., Inskip, C. and Inskip, T. (1998) Pocket Guide to the Birds of Indian Subcontinent. Oxford University Press, Mumbai.
12. Haslem A, Bennett AF. 2008 Birds in agricultural mosaics: the influence of landscape pattern and countryside heterogeneity. *Ecol Appl*.18.185-196.
13. Hussain K.J., Ramesh T., Satpathy K. and Selvanayagam M. (2011). A checklist of birds of Department of Atomic Energy Campus, Kalpakkam, Tamil Nadu. *ZOO's Print.*, 25 (7), 13-20
14. Iahwara Bhat P, Christopher S S, Hosetti BB. 2009. Avifaunal diversity of Anekere Wetland, Karkala, Udupi district, Karnataka, India. 2009. 30(6):1059-62. *J Environ Biol.* 2009 Nov;30(6):1059-62.
15. Rahmani, A.R., Islam, M.Z. and Kasambe, R.M. (2016) Important Bird and Biodiversity Areas in India Priority sites for conservation Second Edition: Revised and Updated Volume
16. Islam M Z and Rahmani A R. Important Bird Areas in India: Priority for Conservation (Revised and updated). Bombay Natural History Society, Indian Bird Conservation Network, Royal Society for the Protection of Birds and BirdLife International (U.K.). Pp. 1992
17. Jayson E A and Mathew D N. 2000. Diversity and species-abundance distribution of birds in the tropical forests of Silent Valley, Kerala. *Journal of the Bombay Natural History Society*. 97(3):390-400
18. Judd, W.S., Campbell, C.S., Kellogg, E.A. and Stevens, P.F. (1999) Plant Systematics—A Phylogenetic Approach. Sinauer Associates, Sunderland
19. Lee PF, Ding TS, Hsu FH, Geng S. 2004 Breeding bird species richness in Taiwan: distribution on gradients of elevation, primary productivity and urbanization. *Journal of Biogeography*. 31(2):307–314.
20. Main streaming conservation of migratory soaring birds into key productive sector along the rift valley/ red sea flyway, bird identification manual, ministry of environment.

21. Malik, D.S. & Joshi N. (2013). Habitat selection pattern of migratory avifauna in relation to nutrients in Asan wetland at Doon valley(Garhwal Himalaya), India. International Journal of Recent ScientificResearch, 4(10), 1470–1475.
22. Manjunatha and Joshi, B. (2012). Avifaunal diversity in Gulbarga region, north Karnatak. Recent Research in Science and Technology 4(7):27-34.
23. Rajashekhar S, Venkatesh MG.2010 The diversity and abundance of water birdsin lakes of Bangalore city, Karnataka, India. Biosystematics. 4(2): 63-73
24. Rapoport, E. H. 1993. The process of plant colonization in small settlements and large cities. In *Humans as components of ecosystems*, ed. M. J. McDonnell and S. T. A. Pickett, 190-207. New York: Springer-Verlag
25. Sinha RK, Dubey M. India as a Megadiversity Nation. Archives of enviro27.Kumar A, Sati JP, et al. Handbook on Indian Wetland Birds and their
26. Sonika K, Amita K, kumar .A S. Maheshwari 2015 Avifaunal Diversity of Tikamgarh District, Madhya Pradesh, India, Discovery Nature, 9(20), , 20-32
27. Stevenson, T. and Fanshawe, J. (2002) Field Guide to the Birds of East Africa: Kenya, Tanzania, Uganda, Rwanda and Burundi. 1st Edition, Princeton University Press, Princeton.
28. Stewart, R.E., Jr. 1996. "Wetlands as Bird Habitat," in J.D. Fretwell, J.S. Williams, and P.J. Redman (eds.) National Water Summary on Wetland Resources, USGS Water-Supply Paper 2425. U.S. Dept. of the Interior, U.S. Geological Survey, Wash., DC, pp. 49-56.